

Acetylcysteine for mucolysis

Newborn use only

2020

Alert	Also known as N-acetylcysteine (NAC). Refer to acetylcysteine intravenous for paracetamol overdose. Safety data for acetylcysteine as a mucolytic agent in newborn infants is limited and the dosage recommendation was on the basis of consensus. Injection preparations are safe to use as oral preparation.
Indication	Meconium ileus secondary to cystic fibrosis Meconium-related ileus of preterm infants Distal intestinal obstruction secondary to cystic fibrosis Gastric or intestinal milk curd obstruction (lactobezoar)
Action	Reduces the viscosity of mucus by cleaving disulphide bonds in the mucoprotein
Drug type	Mucolytic agent
Trade name	DBL Acetylcysteine Injection Concentrate Acetadote Concentrated Injection (solution for infusion) Acetylcysteine-Link Concentrate for infusion
Presentation	DBL Acetylcysteine Injection Concentrate 20%; 200 mg/mL ampoule Acetadote Concentrated Injection (Solution for infusion) 20%; 200 mg/mL vial Acetylcysteine-Link Concentrate for infusion 20%; 200 mg/mL ampoule
Dosage	Intragastric Meconium ileus: 400 mg/dose (4 mL/dose of acetylcysteine 10%) (range 100–500 mg/dose, 1–5 mL/dose of acetylcysteine 10%) 6–8 hourly ⁵ . Acetylcysteine 10% = 100 mg/mL Distal intestinal obstruction secondary to CF in 1–3-month old: 400 mg/dose (8 mL of acetylcysteine 5% (50 mg/mL) daily ²⁹ . Acetylcysteine 5% = 50 mg/mL Rectal enema OR via distal intestinal stoma 40–200 mg/dose of acetylcysteine 4% (1–5 mL of acetylcysteine 4%) 6–8 hourly. Acetylcysteine 4% = 40 mg/mL 50–100 mg/dose of acetylcysteine 1% (5–10 mL/kg of acetylcysteine 1%) 6–8 hourly has also been reported. Acetylcysteine 1% = 10 mg/mL
Dose adjustments – special scenarios	No information.
Maximum dose	400 mg/dose orally
Total cumulative dose	
Route	Oral via gastric tube Rectal Distal intestinal stoma or via T-tube ileostomy Irrigation through Replogle tube
Preparation	Intragastric preparation ^{4,5} Acetylcysteine 10%: Dilute 5 mL of acetylcysteine 20% (200 mg/mL) with 5 mL of glucose 5% or sodium chloride 0.9%* to make a final volume of 10 mL with a concentration of acetylcysteine 10% (100 mg/mL) *For Acetadote Concentrated Injection – use glucose 5% only Rectal or stoma administration ^{22,23} Acetylcysteine 4%: Dilute 1 mL of acetylcysteine 20% (200 mg/mL) with 4 mL of glucose 5% or sodium chloride 0.9%* to make a final volume of 5 mL with a concentration of acetylcysteine 4% (40 mg/mL) *For Acetadote Concentrated Injection – use glucose 5% only.
Administration	Intragastric/rectal/stoma: Administer slowly Irrigation via Replogle tube: As a continuous irrigation with suction applied
Monitoring	Cardiorespiratory, serum electrolytes, liver function
Contraindications	Hypersensitivity to acetylcysteine or any component of the preparation
Precautions	Do not use if intestinal perforation is suspected Abnormal liver and/or renal function

Acetylcysteine for mucolysis

Newborn use only

2020

	Caution in asthma and bronchospasm Acetylcysteine is not compatible with rubber and some metals, particularly, iron, copper and nickel. Can be used with silicone and plastic.
Drug interactions	Glyceryl trinitrate: Increased risk of hypotension
Adverse reactions	Flushing, pruritus and urticarial reactions reported. Hypersensitivity reactions, vomiting, nausea, hypernatremia, hepatotoxicity, mucosal injury and haemorrhage.
Compatibility	Fluids: DBL Acetylcysteine Injection Concentrate and Acetylcysteine-Link Concentrate: Glucose 5%, sodium chloride 0.9% Acetadote Concentrated Injection: Glucose 5%
Incompatibility	Fluids: No information
Stability	Extemporaneously prepared solutions of 1% and 10% acetylcysteine diluted with sodium chloride 0.9% are stable for up to 60 days when stored in plastic amber bottles at room temperature. ²⁴ Acetadote Concentrated Injection is stable for 24 hours at 2 to 8°C after dilution.
Storage	Injection: Store at room temperature. Protect from light.
Excipients	DBL Acetylcysteine Injection Concentrate and Acetylcysteine-Link Concentrate for infusion: Disodium edetate, sodium hydroxide, water for injections. Acetadote Concentrated Injection (solution for infusion): Water for injections and sodium hydroxide for pH adjustment. ²⁶⁻²⁸
Special comments	
Evidence	<p>Acetylcysteine is used for acetaminophen poisoning and as adjuvant therapy in respiratory conditions in paediatric patients. It also exhibits a mucolytic action through its free sulfhydryl group, which opens up the disulfide bonds in mucoprotein to decrease mucus viscosity. For treatment of meconium ileus (MI) or distal intestinal obstruction syndrome (DIOS), acetylcysteine can be given enterally by mouth or by feeding tube. When administered orally, acetylcysteine has been associated with uncommon adverse effects including nausea, vomiting, diarrhoea, dyspepsia and skin rashes. Repeated administration by enema has been associated with hypernatraemia and liver injury.</p> <p>Efficacy</p> <p>Meconium ileus secondary to cystic fibrosis</p> <p>There are no trials of oral or rectal acetylcysteine for meconium ileus secondary to cystic fibrosis. Case series of use of rectal acetylcysteine in uncomplicated meconium ileus include a report of 8 infants were given an acetylcysteine 20%, only 2 were successful in relieving obstruction compared to 9 of 13 relieved with diatrizoic acid (Gastrografin) enema [1]; acetylcysteine and pancreatic enzyme irrigation of a T-tube ileostomy has been reported in two case series with resolution of meconium ileus in 20 of 23 infants using 5–10 mL of an acetylcysteine 1% solution [2]; and in 6 infants using approximately 10 mL of a acetylcysteine 4% solution via the T-tube into the distal ileum [3].</p> <p>Meconium-related obstruction in preterm infants</p> <p>There are no RCTs of oral or rectal acetylcysteine for meconium-related obstruction in preterm infants. Several case series [4, 5] and controlled studies [6, 7] have reported variable efficacy of acetylcysteine orally or rectally in preterm infants.</p> <p>In a retrospective controlled study [6], 132 preterm infants <1250 g with meconium obstruction were given first-line saline rectal irrigations 5–10 mL/kg every 6 hours and metoclopramide 0.1 mg/kg/dose PO or IV every 6 hours. Infants received either oral acetylcysteine 100 mg/kg (10% solution = 1 mL) every 6 hours (n = 34) or rectal enema (1 mL of acetylcysteine 10% [100 mg] added to 9 mL sodium chloride 0.9%) 5–10 mL/kg every 6 hours (n = 52) or no additional treatment (n = 35). There was a reduction in mean time to resolution of obstruction (12 days oral NAC group; 10 days rectal NAC group; 15 days control group) and full enteral feeding. None of the infants was given a contrast enema. Hospital stay and mortality rate did not differ between groups.</p> <p>A before and after study [7] reported 6 of 99 infants born <1500 g in the before period had a meconium-associated bowel obstruction with 4 perforated and 6 surgically managed, compared to 18</p>

of 42 (43%) diagnosed with meconium-related bowel obstruction in the after period. Twelve of 18 resolved with sodium chloride 0.9% enemas, whilst the other 6 resolved with acetylcysteine 100 mg/mL (dose not reported) through an orogastric tube and ultrasound guided diatrizoic acid (Gastrografin) enemas. None required surgery. No complications arose relating to the conservative treatment nor were there any bowel perforations.

Gastric or intestinal milk curd obstruction (lactobezoar)

There are case reports [8-10] of use of acetylcysteine for both gastric and intestinal obstruction with milk curds (lactobezoar) [11]. Successful treatment of gastric lactobezoar was reported using 10 mg/kg/dose of acetylcysteine 10% diluted with 50 mL of sodium chloride 0.9% administered via nasogastric tube over 30 minutes followed by clamping of the nasogastric tube for two hours and repeated 6 hourly up to 7 doses in total [8-10]. A case series reported two extremely preterm infants with intestinal obstruction secondary to human-milk-fortifier-associated curds treated with nasogastrically instilled acetylcysteine and elimination of human milk fortifier [11]. One resolved without surgery whilst the other infant required operative treatment despite nasogastric acetylcysteine and repeated enemas.

Distal intestinal obstruction syndrome (DIOS) secondary to cystic fibrosis

There are no RCTs of oral or rectal acetylcysteine for DIOS associated with cystic fibrosis. The ESPGHAN Cystic Fibrosis Working Group recommend use of acetylcysteine administered orally has been superseded by diatrizoic acid (Gastrografin) in children with an acute episode of DIOS [12]. Oral osmotic laxatives containing polyethylene glycol (PEG) or lactulose are recommended alternatives when needed for prophylaxis against DIOS. [12] A recent review of acetylcysteine for management of DIOS found administration technique and monitoring parameters are not well defined in current literature and clinical trials are lacking and would be helpful to better define the role of acetylcysteine in distal intestinal obstruction syndrome [13].

Maintenance of patency of Replogle tube

There are no reports of use of acetylcysteine for maintenance of patency of a Replogle tube used for suctioning of the proximal pouch in infants with oesophageal atresia.

Safety

In general, oral and rectal acetylcysteine use has been reported to be well tolerated, although the number of reports is insufficient to determine rates of adverse effects. Reported side effects of acetylcysteine treatment include hepatic derangement associated with use of oral acetylcysteine and acetylcysteine 0.2%/contrast enema in an infant with meconium ileus secondary to cystic fibrosis [14]; a preterm infant who developed hypernatraemia with instillation of a 5% solution through a distal stoma at 2.2 mL/hour [15]. There is also a case report of a 3-year-old infant with DIOS associated with cystic fibrosis, who developed liver injury after oral and rectal administration of acetylcysteine [16]. For paracetamol overdose, acetylcysteine is recommended at a dose of 300 mg/kg (200 mg/kg over 4 hours and 100 mg/kg over 16 hours) given intravenously. Adverse reactions are uncommon except for nausea and vomiting. Rashes, erythema, angioedema and anaphylaxis are uncommon [17]. Doses of acetylcysteine are comparable to those documented for paracetamol overdose and, with repeated administration orally or rectally, may substantially exceed these.

Pharmacokinetics/pharmacodynamics

There are few studies on the pharmacokinetics of acetylcysteine with terminal half-life values of between 2.7 and 5.7 hours reported in adults [18]. In infants, gestational age 24.9–31.0 weeks, 2–11 hours after birth, mean elimination half-life was 11 hours (range 7.8–15.2), plasma clearance 37 mL/kg/h (range 13–62) and volume of distribution 573 mL/kg (range 167–1010 mL/kg). A steady-state concentration of acetylcysteine was reached in 2–3 days during a constant infusion. [19] The oral systemic bioavailability of – varied between 6 and 10% in adult volunteers [20, 21], with first-pass metabolism in the liver limiting systemic concentrations [18].

	<p><i>In vitro</i> experiments on constipated mice and human meconium compared perflubron, surfactant, Tween-80, Gastrografin, Golytely, DNase, acetylcysteine 4%, Viokase and sodium chloride 0.9% [22]. For relieving constipation <i>in vivo</i>, Gastrografin enema was most efficacious. All agents were equally benign to the intestinal mucosa. <i>In vitro</i>, only acetylcysteine 4% and perflubron were less effective at decreasing meconium viscosity than sodium chloride 0.9%, with acetylcysteine 4% producing a 69% decrease in viscosity on immediate inspection, but 99% by 6 hours.</p> <p>In another study in 48 puppies aged from 3 days to 3 months, acetylcysteine solutions from 2–20% were compared to sodium chloride 0.9%. Extensive fluid shifts with hyperaemia and multiple haemorrhages of the bowel mucosa were found after exposure to acetylcysteine 20%, with degree of injury varying inversely with the age and size. The severity of the mucosal injury was less with lower concentrations and there were no adverse events reported with the 4% solution. [23]</p> <p>Stability Acetylcysteine 1% (10 mg/mL) and 10% (100 mg/mL) solutions prepared with sodium chloride 0.9% and placed in 2 ounce, amber plastic prescription bottles and stored at 20–25°C were stable with 90% of the initial concentration of acetylcysteine retained in both formulations for 60 days. Neither solution was stable at day 90. [24]</p>
<p>Practice points</p>	<p>For enemas, solutions of acetylcysteine should not exceed 4% to avoid mucosal injury and hypernatraemic dehydration.</p> <p>Monitor electrolytes and liver function tests particularly with repeated administration.</p> <p>There are insufficient data to determine the safety and efficacy of acetylcysteine via nasogastric tube or enema for meconium ileus of prematurity and gastric and intestinal milk curd obstruction (lactobezoar) in preterm infants, particularly in respect to other approaches and agents. [LOE IV, GOR D]</p> <p>Acetylcysteine T-tube ileostomy irrigation has been used for infants with meconium ileus associated with cystic fibrosis. [LOE IV GOR D]</p> <p>The ESPGHAN Cystic Fibrosis Working Group recommend that use of acetylcysteine administered orally has been superseded by diatrizoic acid (Gastrografin) in children with an acute episode of distal intestinal obstruction [12]. Oral osmotic laxatives containing polyethylene glycol (PEG) or lactulose are recommended alternatives when needed for prophylaxis against DIOS. [12] [LOE IV GOR D]</p> <p>There are no published reports of use of acetylcysteine for irrigation of the upper pouch in infants with oesophageal atresia.</p> <p>Irrigation of upper oesophageal pouch in tracheo-oesophageal fistula – no reported evidence. For refractory cases with thick secretions not responding to sodium chloride 0.9% irrigation – subject to surgeon’s approval – 5 mL/hour of acetylcysteine 4% through Replogle tube.</p> <p>Preparation of acetylcysteine 4%: Dilute 20mL of acetylcysteine 20% (200 mg/mL) with 80 mL of glucose 5% or sodium chloride 0.9%* to make a final volume of 100 mL with a concentration of acetylcysteine 4% (40 mg/mL)</p> <p>*For Acetadote Concentrated Injection – use glucose 5% only.</p>
<p>References</p>	<ol style="list-style-type: none"> 1. Garza-Cox S, Keeney SE, Angel CA, Thompson LL, Swischuk LE. Meconium obstruction in the very low birth weight premature infant. <i>Pediatrics</i>. 2004;114:285-90. 2. Mak GZ, Harberg FJ, Hiatt P, Deaton A, Calhoon R, Brandt ML. T-tube ileostomy for meconium ileus: four decades of experience. <i>J Pediatr Surg</i>. 2000;35:349-52. 3. Nguyen LT, Youssef S, Guttman FM, Laberge JM, Albert D, Doody D. Meconium ileus: is a stoma necessary? <i>J Pediatr Surg</i>. 1986;21:766-8. 4. Noblett HR. Treatment of uncomplicated meconium ileus by gastrografin enema: a preliminary report. <i>J Pediatr Surg</i>. 1969;4:190-7. 5. Emil S, Nguyen T, Sills J, Padilla G. Meconium obstruction in extremely low-birth-weight neonates: guidelines for diagnosis and management. <i>J Pediatr Surg</i>. 2004;39:731-7. 6. Kadioglu Simsek G, Arayici S, Buyuktiryaki M, Okur N, Kanmaz Kutman G, Suna Oguz S. Oral N-Acetyl Cysteine for Meconium Ileus of Preterm Infants. <i>Gynecol Obstet Reprod Med</i>. 2019;25:169-73.

7. Solaz-Garcia AJ, Segovia-Navarro L, Rodriguez de Dios-Benlloch JL, Benavent-Taengua L, Castilla-Rodriguez DY, Company-Morena MA. Prevention of meconium obstruction in very low birth weight preterm infants. *Enferm Intensiva*. 2019;30:72-7.
8. Bajorek S, Basaldua R, McGoogan K, Miller C, Sussman CB. Neonatal gastric lactobezoar: management with N-acetylcysteine. *Case Rep Pediatr*. 2012;2012:412412.
9. Savva DA, Crist M, Lardieri A. N-Acetylcysteine for Gastric Lactobezoars in a 1-Month-Old. *J*. 2019;24:247-50.
10. De La Llana SA, Hanquinet S, Pfister RE. Gastric lactobezoar in a Zidovudin treated ELBW infant. *Swiss Medical Weekly*. 2013;197:215.
11. Stanger J, Zwicker K, Albersheim S, Murphy IJ. Human milk fortifier: An occult cause of bowel obstruction in extremely premature neonates. *J Pediatr Surg*. 2014;49:724-6.
12. Colombo C, Ellemunter H, Houwen R, Munck A, Taylor C, Wilschanski M. Guidelines for the diagnosis and management of distal intestinal obstruction syndrome in cystic fibrosis patients. *Journal of Cystic Fibrosis*. 2011;10:S24-S8.
13. Schauble AL, Bisaccia EK, Lee G, Nasr SZ. N-acetylcysteine for Management of Distal Intestinal Obstruction Syndrome. *J*. 2019;24:390-7.
14. Cooke A, Deshpande AV, Wong CK, Cohen R. Hepatic derangement following N-Acetylcysteine enemas in an infant with cystic fibrosis. *J Paediatr Child Health*. 2008;44:673-5.
15. Langer JC, Paes BM, Gray S. Hybernemia associated with N-acetylcysteine therapy for meconium ileus in a premature infant. *CMAJ Canadian Medical Association Journal*. 1990;143:202-3.
16. Bailey DJ, Andres JM. Liver injury after oral and rectal administration of N-acetylcysteine for meconium ileus equivalent in a patient with cystic fibrosis. *Pediatrics*. 1987;79:281-2.
17. Prescott L. Oral or intravenous N-acetylcysteine for acetaminophen poisoning? *Annals of emergency medicine*. 2005;45:409-13.
18. Bateman DN, Dear JW. Acetylcysteine in paracetamol poisoning: A perspective of 45 years of use. *Toxicology Research*. 2019;8:489-98.
19. Ahola T, Fellman V, Laaksonen R, Laitila J, Lapatto R, Neuvonen PJ, Raivio KO. Pharmacokinetics of intravenous N-acetylcysteine in pre-term new-born infants. *Eur J Clin Pharmacol*. 1999;55:645-50.
20. Olsson B, Johansson M, Gabrielsson J, Bolme P. Pharmacokinetics and bioavailability of reduced and oxidized N-acetylcysteine. *European Journal of Clinical Pharmacology*. 1988;34:77-82.
21. Borgstrom L, Kagedal B, Paulsen O. Pharmacokinetics of N-acetylcysteine in man. *European Journal of Clinical Pharmacology*. 1986;31:217-22.
22. Burke MS, Ragi JM, Karamanoukian HL, Kotter M, Brisseau GF, Borowitz DS, Ryan ME, Irish MS, Glick PL. New strategies in nonoperative management of meconium ileus. *J Pediatr Surg*. 2002;37:760-4.
23. Shaw A. Safety of N-acetylcysteine in treatment of meconium obstruction of the newborn. *J Pediatr Surg*. 1969;4:119-25.
24. Fohl AL, Johnson CE, Cober MP. Stability of extemporaneously prepared acetylcysteine 1% and 10% solutions for treatment of meconium ileus. *Am J Health-Syst Pharm*. 2011;68:69-72.
25. Society of Hospital Pharmacists of Australia, Australian Injectable Drugs Handbook, 8th Edition 2020. Accessed 26/02/2020.
26. Product Information. Acetadote Concentrated Injection (Solution for infusion). MIMS online. Accessed 26/02/2020.
27. Product Information. Acetylcysteine-Link Concentrate for infusion. MIMS online. Accessed 26/02/2020.
28. Product Information. DBL Acetylcysteine Injection Concentrate. MIMS online. Accessed 26/02/2020.
29. Cystic fibrosis manual – Children’s Hospital at Westmead practice guideline. Accessed on 6 March 2020.

Acetylcysteine for mucolysis

Newborn use only

2020

VERSION/NUMBER:	DATE
Original: 1.0	26/03/2020
REVIEW	26/03/2025
Approval by: JHCHCQ&PCC	Approval date: 27/05/2020

Authors Contribution

Original author/s	Gayatri Panaboke, Himanshu Popat, Srinivas Bolisetty
Evidence Review	David Osborn
Expert review	Gordon Thomas, Bruce Currie, Ashish Jiwane
Nursing Review	Eszter Jozsa
Pharmacy Review	Cindy Chen, Michelle Jenkins, Carmen Burman
ANMF Group contributors	Nilkant Phad
Final editing and review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty