

Alert	The Antimicrobial Stewardship Team recommends this drug is listed under the following category: Unrestricted.																										
Indication	Directed treatment of infections caused by susceptible gram positive (including <i>Streptococcus</i> species, <i>Enterococcus faecalis</i> and <i>Listeria monocytogenes</i>) and susceptible gram negative bacteria (some strains of <i>Escherichia coli</i> , many strains of <i>Haemophilus influenzae</i> , <i>Neisseria meningitidis</i> , <i>Proteus mirabilis</i> and <i>Salmonellae</i>). Empiric treatment of suspected early onset sepsis including meningitis, with an aminoglycoside.																										
Action	Bactericidal - inhibits the synthesis of the bacterial cell wall. Ampicillin is hydrolysed by beta-lactamases and therefore not effective against penicillinase producing bacteria.																										
Drug Type	Antibacterial - Penicillin																										
Trade Name	Ampicyn, Austrapen, Ibimicyn																										
Presentation	Ampicillin 500 mg vial Ampicillin 1000 mg vial																										
Dosage / Interval	<p>Standard infections: 50 mg/kg/dose. Dosing interval as per table below</p> <p>Meningitis: 100 mg/kg/dose. Dosing interval as per table below</p> <table border="1"> <thead> <tr> <th colspan="2">Method</th> <th rowspan="2">Interval</th> </tr> <tr> <th>Corrected Gestational Age/Postmenstrual Age</th> <th>Postnatal Age</th> </tr> </thead> <tbody> <tr> <td>< 30⁺⁰ weeks</td> <td>0–28 days</td> <td>12 hourly</td> </tr> <tr> <td>< 30⁺⁰ weeks</td> <td>29+ days</td> <td>8 hourly</td> </tr> <tr> <td>30⁺⁰–36⁺⁶ weeks</td> <td>0–14 days</td> <td>12 hourly</td> </tr> <tr> <td>30⁺⁰–36⁺⁶ weeks</td> <td>15+ days</td> <td>8 hourly</td> </tr> <tr> <td>37⁺⁰–44⁺⁶ weeks</td> <td>0–7 days</td> <td>12 hourly</td> </tr> <tr> <td>37⁺⁰–44⁺⁶ weeks</td> <td>8+ days</td> <td>8 hourly</td> </tr> <tr> <td>≥ 45⁺⁰ weeks</td> <td>0+ days</td> <td>6 hourly</td> </tr> </tbody> </table>	Method		Interval	Corrected Gestational Age/Postmenstrual Age	Postnatal Age	< 30 ⁺⁰ weeks	0–28 days	12 hourly	< 30 ⁺⁰ weeks	29+ days	8 hourly	30 ⁺⁰ –36 ⁺⁶ weeks	0–14 days	12 hourly	30 ⁺⁰ –36 ⁺⁶ weeks	15+ days	8 hourly	37 ⁺⁰ –44 ⁺⁶ weeks	0–7 days	12 hourly	37 ⁺⁰ –44 ⁺⁶ weeks	8+ days	8 hourly	≥ 45 ⁺⁰ weeks	0+ days	6 hourly
Method		Interval																									
Corrected Gestational Age/Postmenstrual Age	Postnatal Age																										
< 30 ⁺⁰ weeks	0–28 days	12 hourly																									
< 30 ⁺⁰ weeks	29+ days	8 hourly																									
30 ⁺⁰ –36 ⁺⁶ weeks	0–14 days	12 hourly																									
30 ⁺⁰ –36 ⁺⁶ weeks	15+ days	8 hourly																									
37 ⁺⁰ –44 ⁺⁶ weeks	0–7 days	12 hourly																									
37 ⁺⁰ –44 ⁺⁶ weeks	8+ days	8 hourly																									
≥ 45 ⁺⁰ weeks	0+ days	6 hourly																									
Maximum Daily Dose	400 mg/kg/day																										
Route	IV IM (only if IV route not possible as intramuscular route is painful)																										
Preparation/Dilution	<p>IV: Add 4.7 mL of water for injection to the 500 mg vial for reconstitution to make 100 mg/mL solution OR Add 9.3 mL of water for injection to the 1 g vial for reconstitution to make 100 mg/mL solution. 100 mg/mL can be infused directly, but if desired and fluid balance allows, can be</p> <p>FURTHER DILUTED:</p> <ul style="list-style-type: none"> - Draw up 5 mL (500 mg of ampicillin) of solution and add 5 mL sodium chloride 0.9% to make a final volume of 10mL with a concentration of 50 mg/mL solution OR - Draw up 3 mL (300 mg of ampicillin) of solution and add 7 mL sodium chloride 0.9% to make a final volume of 10mL with a concentration of 30 mg/mL solution <p>IM: Add 1.7 mL of water for injection to the 500 mg vial for reconstitution to make 250 mg/mL solution.</p>																										
Administration	IV: Infusion over 5–10 minutes into the proximal cannula site with a maximum rate of 100 mg/minute. Separate from aminoglycosides by clearing the lines with a flush as ampicillin inactivates them. Higher doses should be diluted to 30 mg/mL and infused over 30 minutes.																										
Monitoring	Plasma concentrations not usually required; however may be useful for infections caused by bacteria with high Minimum Inhibitory Concentration (MIC).																										
Contraindications	Hypersensitivity reactions can occur in ampicillin-treated infants younger than 6 months of age but are rarely reported in neonates.																										

Precautions	Hypersensitivity to penicillin derivatives. In renal impairment the excretion of ampicillin will be delayed. In infants with severe renal impairment it may be necessary to reduce the total daily dose.
Drug Interactions	Aminoglycosides including gentamicin should not be mixed with ampicillin when both drugs are given parenterally as inactivation occurs. Ensure line is adequately flushed between antibiotics.
Adverse Reactions	Allergic reactions – maculopapular or urticarial rash, fever (rare in neonates). Other: Diarrhoea; CNS excitation or seizures with very large doses reported in adults; and prolonged bleeding time with repeated doses.
Compatibility	Fluids: Sodium chloride 0.9%. Y site: Aciclovir, amifostine, anidulafungin, aztreonam, bivalirudin, dexmedetomidine, esmolol, filgrastim, foscarnet, granisetron, heparin sodium, labetalol, linezolid, magnesium sulfate, morphine sulfate, pethidine, potassium chloride, remifentanyl.
Incompatibility	Fluids: Glucose and glucose containing solutions, fat emulsions. Y site: Amino acid solutions, adrenaline hydrochloride, aminoglycosides – amikacin, gentamicin, tobramycin; aminophylline, atropine, buprenorphine, caspofungin, chlorpromazine, clindamycin, dobutamine, dolasetron, dopamine, ergometrine, fluconazole, ganciclovir, haloperidol lactate, hydralazine, ketamine, lincomycin, metoclopramide, midazolam, mycophenolate mofetil, ondansetron, pentamidine, prochlorperazine, promethazine, protamine, sodium bicarbonate, tranexamic acid, verapamil.
Stability	Administer immediately; discard unused portion of reconstituted solution.
Storage	Store below 25°C Protect from light.
Special Comments	Clearance is primarily by the renal route. Clearance increases with increasing gestational age and postnatal age. Serum half-life is longer in premature infants and infants younger than 7 days.
Evidence summary	Refer to full version.
References	Refer to full version.

Original version Date: 08/08/2015	Author: NeoMed Consensus Group
Current Version number: 1.1	Current Version Date: 20/02/2017
Risk Rating: Medium	Due for Review: 26/07/2019
Approval by: JHCH CQ&PCC	Approval Date: 27/05/2020