

Adrenaline (epinephrine) IV infusion

Newborn use only

2020

Alert	1:10,000 (1 mg/10 mL) ampoule is the preferred preparation for adrenaline infusion.												
Indication	Treatment of hypotensive shock with or without myocardial dysfunction.												
Action	<p>Catecholamine with alpha and beta adrenergic actions.</p> <p>Haemodynamic effects are dose dependent:</p> <ul style="list-style-type: none"> • At low doses of 0.01–0.1 microgram/kg/minute primarily stimulates cardiac and vascular beta 1- and beta 2-adrenoreceptors leading to increased inotropy, chronotropy, conduction velocity and peripheral vasodilation. • At doses greater than 0.1 microgram/kg/minute adrenaline also stimulates vascular and cardiac alpha 1-receptors causing vasoconstriction and increased inotropy. The net effects are increases in blood pressure and systemic blood flow caused by the drug-induced increases in systemic vascular resistance (SVR) and cardiac output.¹ 												
Drug type	Inotropic vasopressor.												
Trade name	Aspen Adrenaline 1: 10,000 Adrenaline Acid Tartrate injection; Adrenaline 1:1,000 Adrenalin Acid Tartrate injection.												
Presentation	1 mg/10 mL or 1:10,000 ampoule [100 microgram/mL] 1 mg/mL or 1:1,000 ampoule [1000 microgram/mL]												
Dose	Low dose: 0.05–0.1 microgram/kg/minute High dose: 0.1–1 microgram/kg/minute												
Dose adjustment													
Maximum dose													
Total cumulative dose													
Route	Continuous IV infusion.												
Preparation	<p>Preparation using 1:10,000 (1 mg/10 mL) ampoule</p> <p>LOW CONCENTRATION IV infusion</p> <table border="1" style="width: 100%; border-collapse: collapse; margin-bottom: 10px;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.05 microgram/kg/minute</td> <td>150 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 150 microgram/kg (1.5 mL/kg) of 1:10,000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 3 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.05 microgram/kg/minute.</p> <p>HIGH CONCENTRATION IV infusion</p> <table border="1" style="width: 100%; border-collapse: collapse; margin-bottom: 10px;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.2 microgram/kg/minute</td> <td>600 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 600 microgram/kg (6 mL/kg) of 1:10,000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 12 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.2 microgram/kg/minute.</p> <p>For infants requiring fluid restriction consider: VERY HIGH CONCENTRATION IV infusion*</p> <table border="1" style="width: 100%; border-collapse: collapse; margin-bottom: 10px;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.4 microgram/kg/minute</td> <td>1200 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 1200 microgram/kg (12 mL/kg) of 1:10,000 adrenaline and add glucose 5% ONLY to make a final volume of 50 mL with a concentration of 24 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.4 microgram/kg/minute.</p> <p>*Stability data only available for 5% glucose for very high concentration.</p> <p>Preparation using 1:1,000 (1 mg/mL) ampoule – Occasionally used for infants >4 kg: 1:1000 (1 mg/mL) ampoule is not commonly kept in the NICUs</p>	Infusion Strength	Prescribed amount	1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL	Infusion Strength	Prescribed amount	1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL	Infusion Strength	Prescribed amount	1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL
Infusion Strength	Prescribed amount												
1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL												
Infusion Strength	Prescribed amount												
1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL												
Infusion Strength	Prescribed amount												
1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL												

	<p>LOW CONCENTRATION IV infusion</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.05 microgram/kg/minute</td> <td>150 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 150 microgram/kg (0.15 mL/kg) of 1:1000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 3 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.05 microgram/kg/minute.</p> <p>HIGH CONCENTRATION IV infusion</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.2 microgram/kg/minute</td> <td>600 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 600 microgram/kg (0.6 mL/kg) of 1:1000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 12 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.2 microgram/kg/minute.</p> <p>For infants requiring fluid restriction consider: VERY HIGH CONCENTRATION IV infusion*</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">Infusion Strength</th> <th style="width: 50%;">Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.4 microgram/kg/minute</td> <td>1200 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 1200 microgram/kg (1.2 mL/kg) of 1:1000 adrenaline and add glucose 5% ONLY to make a final volume of 50 mL with a concentration of 24 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.4 microgram/kg/minute.</p> <p>*Stability data only available for 5% glucose for very high concentration.</p>	Infusion Strength	Prescribed amount	1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL	Infusion Strength	Prescribed amount	1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL	Infusion Strength	Prescribed amount	1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL
Infusion Strength	Prescribed amount												
1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL												
Infusion Strength	Prescribed amount												
1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL												
Infusion Strength	Prescribed amount												
1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL												
Administration	Continuous IV infusion preferably via dedicated central line. Use with caution via a peripheral line.												
Monitoring	Continuous heart rate, ECG and blood pressure monitoring preferable. Assess urine output and peripheral perfusion frequently. Observe IV site closely for blanching and extravasation.												
Contraindications	Arrhythmia and tachyarrhythmia. Cardiovascular disease resulting in arterial narrowing including cerebrovascular disease, coronary artery disease and digital ischaemia. Pheochromocytoma. Thyrotoxicosis. Glaucoma. Known hypersensitivity to sympathomimetic amines.												
Precautions	Ensure adequate circulating blood volume prior to commencement. Potent chronotrope and vasopressor – may cause excessive tachycardia, severe hypertension and ventricular arrhythmias. May cause lactic acidosis and hyperglycaemia.												
Drug interactions	Hypotension may be observed with concurrent use of vasodilators such as glyceryl trinitrate, nitroprusside and calcium channel blockers. Concurrent use of digitalis glycosides may increase the risk of cardiac arrhythmias. Concurrent use of IV phenytoin with adrenaline may result in dose dependent, sudden hypotension and bradycardia.												
Adverse reactions	Tachycardia and arrhythmia. Systemic hypertension especially at higher doses. May cause hypokalaemia. Tissue necrosis at infusion site with extravasation. Digital ischaemia.												
Compatibility	Fluids: Glucose 5%, glucose 10%, Hartmann's, sodium chloride 0.9%. Stability data only available for 5% glucose for very high concentration.												

Adrenaline (epinephrine) IV infusion

Newborn use only

2020

	<p>Y-site: Amino acid solutions. Amiodarone, anidulafungin, atracurium, bivalirudin, caspofungin, cisatracurium, dexmedetomidine, dobutamine, dopamine, ethanol, fentanyl, glyceryl trinitrate, heparin sodium, milrinone, morphine sulfate, pancuronium, potassium chloride, ranitidine, remifentanyl, sodium nitroprusside, tigecycline, tirofiban, vecuronium.</p> <p>No information: Adrenaline HCL is compatible with noradrenaline bitartrate but no stability data is available for Adrenaline acid tartrate and noradrenaline bitartrate</p>
Incompatibility	<p>Fluids: Sodium bicarbonate.</p> <p>Y-site: Aciclovir, aminophylline, ampicillin, atropine, azathioprine, calcium chloride, calcium gluconate, cefalotin, chloramphenicol, digoxin, ergometrine, ganciclovir, hyaluronidase, hydrocortisone sodium succinate, indomethacin, phenobarbitone sodium, sodium bicarbonate, thiopentone, vancomycin.</p>
Stability	Diluted solution: Stable for 24 hours below 25°C.
Storage	Store below 25°C. Protect from light. Discard remainder after use.
Excipients	
Special comments	<p>Preferably administered via "dedicated" line to avoid accidental bolus. Do not use as a side line with maintenance fluids.</p> <p>Discard if exhibiting colour change.</p>
Evidence	Refer to full version.
Practice points	Refer to full version.
References	Refer to full version.

VERSION/NUMBER	DATE
Original 1.1	31/03/2016
Revised 1.2	30/11/2017
Current 2.0	5/01/2021
REVIEW	5/01/2026

Authors Contribution

Original author/s	David Osborn
Evidence Review	David Osborn
Expert review	
Nursing Review	Eszter Jozsa
Pharmacy Review	Jing Xiao, Mariella De Rosa, Ushma Trivedi, Cindy Chen
ANMF Group contributors	Ansar Kunjunju, Srinivas Bolisetty, David Osborn, Eszter Jozsa, Nilkant Phad, Cindy Chen
Final editing and review of the original	Ian Whyte
Electronic version	Mariella De Rosa, Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty