

Chloramphenicol Topical

Newborn use only

2020

Alert	Eye ointment remains in the eye for longer than eye drops and may be the preferred dose form.
Indication	Treatment of acute bacterial conjunctivitis
Action	Bacteriostatic. Acts by inhibition of protein synthesis, interfering with the transfer of activated amino acids from soluble RNA to ribosomes.
Drug type	Broad spectrum antibiotic
Trade name	Chloromycetin, Chlorsig
Presentation	Chloramphenicol eye drops 0.5% contains 5 mg/mL, chloramphenicol eye ointment 1% contains 10mg/g
Dose	<p>Ointment: Apply 3 to 4 times a day in the affected eye and continue for 48 hours after clinical resolution. Severe infection: May need more often at the discretion of the treating team.</p> <p>DROPS: Severe infection: Apply 1 drop every 2 hours in the affected eye for 48 hours and reduce frequency with controlling of infection. Less severe infection: Apply 3 to 4 times a day in the affected eye Continue for 48 hours after clinical resolution</p> <p>Drops and Ointment combination: Apply drops during the day as above Apply ointment once at night Continue for 48 hours after clinical resolution.</p>
Dose adjustment	
Therapeutic hypothermia	Not applicable
ECMO	Not applicable
Renal	Not applicable
Hepatic	Not applicable
Maximum dose	
Total cumulative dose	
Route	TOPICAL ONLY
Preparation	
Administration	<p>Avoid contact between tip of container and infant's eyes.</p> <p>Eye ointment: Hold eye open and administer eye ointment between the lower lid and the eye.</p> <p>Eye drop: After administering eye drop, gently press against the lacrimal duct (inner corner of eye) to reduce systemic absorption. The eye pouch will be full after a single drop. If other eye drop(s) need to be administered, wait 5 minutes between drops.</p>
Monitoring	
Contraindications	History of hypersensitivity to chloramphenicol or any other component of the medication.
Precautions	Family history of blood disorders
Drug interactions	
Adverse reactions	<p>Local irritation e.g. burning, swelling, redness; impaired corneal healing; superinfection; hypersensitivity including sensitisation, urticaria, rash, fever, angioedema, anaphylaxis, blood dyscrasia (rare).</p> <p>Acute hepatitis was reported in an adult following topical chloramphenicol therapy for conjunctivitis.</p> <p>Bone marrow hypoplasia, including aplastic anaemia and death, has been rarely reported following local application of chloramphenicol.</p> <p>Overgrowth of non-susceptible organisms.</p>
Compatibility	No information
Incompatibility	No information
Stability	

Chloramphenicol Topical

Newborn use only

2020

Storage	Eye drop: Store unopened bottle at 2–8°C. Once opened, bottle may be stored at <25°C for 28 days. Protect from light. Eye Ointment: Store below 25°C. Discard 28 days after opening. Protect from light
Excipients	
Special comments	
Evidence	Please see the full version
Practice points	Chloramphenicol topical eye drops may be preferable if infection of blocked lacrimal apparatus is suspected and if the use is intended at home.
References	Please see the full version

VERSION/NUMBER	DATE
Original	21/06/2019
Revised 1.1	02/07/2020
Current 1.1	02/07/2020
REVIEW (5 years)	01/07/2025

Authors Contribution

Original author/s	Srinivas Bolisetty	
Evidence Review	David Osborn	
Expert review	Tony Lai, Brendan McMullan, Alison Kesson	
Nursing Review	Eszter Jozsa	
Pharmacy Review	Cindy Chen, Jing Xiao, Michelle Jenkins	
ANMF Group contributors	Nilkant Phad, Himanshu Popat	
Final editing and review of the original	Ian Whyte	
Electronic version	Cindy Chen, Ian Callander	
Facilitator	Srinivas Bolisetty	