

# Adrenaline (epinephrine) nebulised

## Newborn use only

2021

<b>Alert</b>	Adrenaline <b>1:1000</b> strength should be used for nebulisation.
<b>Indication</b>	Management of post-extubation stridor [evidence for effectiveness is not clear]. [1-4] Initial treatment of outpatients with moderate to severe bronchiolitis. [5] Initial treatment of croup.[6]
<b>Action</b>	Sympathomimetic catecholamine with alpha and beta adrenergic actions. Vasoconstrictor. It also induces relaxation of the bronchial smooth muscle by acting on beta-adrenergic receptors to alleviate wheezing and dyspnoea. The effects of nebulised adrenaline for the treatment of croup lasts for 2–3 hours.
<b>Drug type</b>	Sympathomimetic catecholamine. Inotropic vasopressor.
<b>Trade name</b>	Adrenaline <b>1:1,000</b> Adrenaline Acid Tartrate injection.
<b>Presentation</b>	1 mg/mL or <b>1:1,000</b> ampoule [1000 microgram/mL]
<b>Dose</b>	0.5 mg/kg (0.5 mL/kg of adrenaline 1:1000 ampoule) Dose may be repeated every 60 minutes if required following medical assessment of previous dose effect.
<b>Dose adjustment</b>	Not applicable.
<b>Maximum dose</b>	
<b>Total cumulative dose</b>	
<b>Route</b>	Nebulised
<b>Preparation</b>	<b>Using a 1:1,000 (1000 microgram/1 mL) ampoule</b> Draw up 0.5 mL/kg (0.5 mg/kg) adrenaline and add sodium chloride 0.9% to make a final volume of 4 mL.
<b>Administration</b>	Deliver final volume of 4 mL via nebuliser over 15 minutes. Driving gas as prescribed by medical staff. Set flow rate at 6 L/minute. There will always be dead space that is not available for nebulisation - it is not possible to nebulise to dryness.
<b>Monitoring</b>	Administer under close supervision of medical staff. Cardiorespiratory monitoring including respiratory rate, oxygen saturation, heart rate and blood pressure.
<b>Contraindications</b>	
<b>Precautions</b>	Infants with arrhythmias, hypertension or hyperthyroidism. Infants with dilated or ischaemic cardiac disease (relative). Crosscheck correct adrenaline strength ampoule used. Do not use if the injection is pink or brown or contains a precipitate.
<b>Drug interactions</b>	No information.
<b>Adverse reactions</b>	Tachycardia and arrhythmia. Systemic hypertension.
<b>Compatibility</b>	Fluids: Sodium chloride 0.9% Drugs: No information. Not to be mixed with other drugs in the same nebulisation chamber.
<b>Incompatibility</b>	Fluids and drugs: No information.
<b>Stability</b>	Discard remainder after use.
<b>Storage</b>	Store below 25°C. Protect from light. Do not refrigerate or freeze.
<b>Excipients</b>	Tartaric acid, sodium metabisulfite, sodium chloride and water for injections.
<b>Special comments</b>	
<b>Evidence</b>	<b>Efficacy:</b> <b>Nebulised racemic adrenaline for extubation of newborn infants:</b> There are no trials proving the efficacy of nebulised adrenaline compared to placebo or intravenous dexamethasone for post extubation stridor. [1-4]

	<p><b>Treatment and prevention of bronchiolitis in newborns and infants:</b> Nebulised adrenaline decreases hospitalisations in patients presenting to ER. There is no evidence to support the use of epinephrine for inpatients. [5, 8] (LOE I, GOR A)</p> <p><b>Treatment of children with croup:</b> Nebulised epinephrine is associated with clinically and statistically significant transient reduction of symptoms of croup 30 minutes post-treatment. [6] (LOE I, GOR A) Evidence does not favour racemic epinephrine or L-epinephrine, or IPPB over simple nebulization. (LOE II, GOR B)</p> <p><b>Safety:</b> Nebulised adrenaline is associated with increased heart rate and blood pressure. [2, 8]</p> <p><b>Pharmacokinetics:</b> Not reported for nebuliser use in newborns or children. No difference in plasma adrenaline levels in asymptomatic children with history of anaphylaxis given adrenaline inhaler (10-20 activations) versus children given a placebo.[9]</p>
<b>Practice points</b>	
<b>References</b>	<ol style="list-style-type: none"> <li>1. Cesar RG, de Carvalho WB. L-epinephrine and dexamethasone in postextubation airway obstruction: a prospective, randomized, double-blind placebo-controlled study. <i>International journal of pediatric otorhinolaryngology</i>. 2009;73:1639-43.</li> <li>2. da Silva PS, Fonseca MC, Iglesias SB, Junior EL, de Aguiar VE, de Carvalho WB. Nebulized 0.5, 2.5 and 5 ml L-epinephrine for post-extubation stridor in children: a prospective, randomized, double-blind clinical trial. <i>Intensive care medicine</i>. 2012;38:286-93.</li> <li>3. Davies MW, Davis PG. Nebulized racemic epinephrine for extubation of newborn infants. <i>The Cochrane database of systematic reviews</i>. 2002:CD000506.</li> <li>4. Preutthipan A, Poomthavorn P, Sumanapisan A, Chinrat B, Thasuntia S, Plitponkarnpim A, Chantarojanasiri T. A prospective, randomized double-blind study in children comparing two doses of nebulized L-epinephrine in postintubation croup. <i>Journal of the Medical Association of Thailand = Chotmaihet thangphaet</i>. 2005;88:508-12.</li> <li>5. Baraldi E, Lanari M, Manzoni P, Rossi GA, Vandini S, Rimini A, Romagnoli C, Colonna P, Biondi A, Biban P, Chiamenti G, Bernardini R, Picca M, Cappa M, Magazzu G, Catassi C, Urbino AF, Memo L, Donzelli G, Minetti C, Paravati F, Di Mauro G, Festini F, Esposito S, Corsello G. Inter-society consensus document on treatment and prevention of bronchiolitis in newborns and infants. <i>Italian journal of pediatrics</i>. 2014;40:65.</li> <li>6. Bjornson C, Russell K, Vandermeer B, Klassen TP, Johnson DW. Nebulized epinephrine for croup in children. <i>The Cochrane database of systematic reviews</i>. 2013;10:CD006619.</li> <li>7. Muraro A, Roberts G, Worm M, Bilo MB, Brockow K, Fernandez Rivas M, Santos AF, Zolkipli ZQ, Bellou A, Beyer K, Bindslev-Jensen C, Cardona V, Clark AT, Demoly P, Dubois AE, DunnGalvin A, Eigenmann P, Halcken S, Harada L, Lack G, Jutel M, Niggemann B, Rueff F, Timmermans F, Vlieg-Boerstra BJ, Werfel T, Dhimi S, Panesar S, Akdis CA, Sheikh A, Allergy EF, Anaphylaxis Guidelines G. Anaphylaxis: guidelines from the European Academy of Allergy and Clinical Immunology. <i>Allergy</i>. 2014;69:1026-45.</li> <li>8. Hartling L, Bialy LM, Vandermeer B, Tjosvold L, Johnson DW, Plint AC, Klassen TP, Patel H, Fernandes RM. Epinephrine for bronchiolitis. <i>The Cochrane database of systematic reviews</i>. 2011:CD003123.</li> <li>9. Simons FE, Gu X, Johnston LM, Simons KJ. Can epinephrine inhalations be substituted for epinephrine injection in children at risk for systemic anaphylaxis? <i>Pediatrics</i>. 2000;106:1040-4.</li> <li>10. Australian Injectable Drugs Handbook, 6th Edition, Society of Hospital Pharmacists of Australia 2014</li> </ol>

VERSION/NUMBER	DATE
Original 1.0	5/05/2016
Current 2.0	22/04/2021
REVIEW	22/04/2026

**Authors Contribution**

Original author/s	David Osborn, Srinivas Bolisetty
Evidence Review	David Osborn
Expert review	
Nursing Review	Eszter Jozsa
Pharmacy Review	Jessica Mehegan
ANMF Group contributors	Ansar Kunjunju, Nilkant Phad, Bhavesh Mehta, John Sinn, Michelle Jenkins, Thao Tran, Helen Huynh
Final editing and review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty